

BIDHAN CHANDRA KRISHI VISWAVIDYALAYA
MEDIUM-RANGE WEATHER FORECAST AND AGRO-ADVISORY
(Agromet Advisory Services, IMD, MoES)

Directorate of Research,
B.C.K.V., Kalyani.



E-mail : aaskalvani@gmail.com
bhowmickmahasweta2012@gmail.com

From Kalyani (Nadia, WB)

Weather Forecast for Hooghly District

FORECAST PERIOD: : 15th September to 19th September, 2018

Weather forecast for next five days

According to SPI data sent by IMD the weather of New Alluvial Zone of West Bengal will be Moderately Dry and Mildly Wet in the coming week.

- Mainly cloudy sky is expected during next few days .
- Light rainfall is expected during next few days and total will be 23 mm.
- Wind flow mainly from S, SW, SE,NW, N & W direction with 1.2-2.1 km/hr. speed during next 5 days is expected.
- Maximum relative humidity is expected to be around 69%-90% and minimum relative humidity is expected to be around 47% - 56%.
- Maximum temperature is expected to be around 34°C-35°C and minimum temperature is expected to be around 26 °C -27°C .

Agrometeorological Advisory

According to NDVI data sent by IMD agricultural vigour of New Alluvial Zone of West Bengal will be good in the coming week.

Crop	Stage	Advisory
Corn	Sowing	<i>Farmers are advised for sowing of sweet corn & baby corn in raised bed. Treat the seed with Carbendazim 50WP @ 2 gm/kg seed or Mancozeb 45WP @ 2.5gm/kg seed or Trichoderma spp. @ 5 gm/kg. seed.</i>
Aman	Vegetative	At this time paddy crop is in vegetative stage. Farmers are advised to monitor crop regularly, if population of leaf folder or stem borer is

		above ETL than broadcast of Cartap 4% granules or carbofuran 3% G @ 10 kg/acre or chlorpyrifos 20 EC @ 2 ml/ litre of water for effective control is recommended. In present weather condition, constant monitoring for brown plant hopper in paddy field is advised.. If insect population is above ETL spray of Imidacloprid 17.8 % SC@ 1.0 ml / 3 lit of water or is recommended.Spray Tricyclozole 75%WP @ 1.5gm/L or Propiconazole 25%EC @ Hexaconazole 5EC @ 1.5ml/L or Carbendazim 50WP @ 1.0gm/L , if there is sheath blight disease appearance in the field.
Bhindi, Brinjal , Chilly	Vegetative	Constant monitoring of ladyfinger, brinjal and chilli crop against attack of mite and jassid is advised. To control mite, spraying of phosmite @ 2 ml per litre of water is advised while spraying of Dimethoate 30 EC @ 2 ml/liter of water is advised against jassid when sky remain clear.
Cabbage, Caulifl ower	Vegetative	Farmers are advised to monitor borer insect. If insect population is above ETL than, spraying of Spinosad 1.0 ml/4 litre of water is advised when sky is clear against fruit borer, diamond back moth etc.
Chrysant hemum & Marigold	Sowing	Farmers are advised to prepare marigold & chrysanthemum nursery in net house. Proper drainage should be done in the fields. Apply <i>Trichoderma</i> spp. along with organic matter@ 5 gm/kg. of organic matter in the nursery bed soil.
