



PurbaMedinipur

South 24 Parganas

(Period to 19th January to 23rd January, 2019)

Issued jointly by GKMS, Kakdwip, BCKV and IMD



Bulletin No.:06/2019

Date: 18th January, 2018

Weather for last three days (15th January to 17th January, 2019)

	15/01/18	16/01/18	17/01/19	
Rainfall (mm)	0.0	0.0	0.0	
Maximum Temperature (°C)	23.0	25.0	26.0	
Minimum Temperature (°C)	11.0	13.0	14.0	
Relative Humidity (%)	60	80	95	
Wind Speed (Km ph)				

Weather for the preceding week (11th January to 17th January, 2018)

Rainfall (mm)	0.0
Maximum Temperature (°C)	21.0-27.0
Minimum Temperature (°C)	10.0-15.0
Relative Humidity (%)	52-95
Wind Speed (Km ph)	

Weather forecast for next five days (19th January to 23rd January, 2019)

- No rainfall is predicted in the next five days.
- Sky will be clear in the next five days.
- Wind speed will be 5.9-8.7 km/sec. and the predominant wind direction will be South-Westerly to Northerly.
- Maximum temperature is expected to be around 26.0^oC -29.0^oC and minimum temperature is likely to be around 12.0^oC-15.0^oC.
- Maximum and minimum relative humidity will be in the range of 51% -55% and 24% -28%.

Weather based Agro-Advisories

Crop	Stage	Disease pest	Advisory
Boro paddy	Seedling (>30days old)		--Prepare main field by proper ploughing and removing all the weeds and transplanting start within 2-3 days --Apply basal fertilizer viz.urea@9kg/bigha , SSp@54kg/bigha and MOP@11kg/bigha --Apply Phorate granule @ 40g/decimal or Carbofuran 3G @120g/decimal over seed bed , 7 days before transplanting
	Seedlings in seed bed	Yellowing due to morning fog	--For preventing fog make shade by thin polythene --For yellowing of seed lings apply chelated Zinc or Zinc EDTA or SAF @1g/liter of water after complete drying of seed bed
Potato	Tuber formation	Late blight	--Where plants found already attacked apply a mixture of Thiamethoxin and Mancozeb@ 3g/liter of water or a mixture of Diomithomorf and Mancozeb @3g/liter of water --Apply all the medicine alternatively --Wet the plant fully by insecticide --Stop irrigation and urea application
Lathyrus	Start flowering	Yellowing of plant due to fog	--Apply Diethane-M-45 @2.5g/liter of water --Apply 2% DAP for better flowering
Fishery			--Apply less amount of floating food only day time --Clean the surrounding for direct sunlight --Stirring the fishes with net must be done per week --Apply 7-10 kg lime mixed with water per bigha pond area during morning time --Start to sell Ruhi, Katla, Mrigel etc. carp

**Good Agriculture vigour was found from NDVI data and moderate wet condition was found from SPI data in the Southern Part of West Bengal*

