

BIDHAN CHANDRA KRISHI VISWAVIDYALAYA
MEDIUM-RANGE WEATHER FORECAST AND AGRO-ADVISORY
(Agromet Advisory Services, IMD, MoES)

Directorate of Research,
B.C.K.V., Kalyani.



Tel : (033) 2582-8721
E-mail : aaskalyani@rediffmail.com

From Kalyani (Nadia, WB)

Weather Forecast for Burdwan District
FORECAST PERIOD: 26th July to 30th July, 2017

Weather forecast for next five days

- Mainly cloudy sky on all of the days is expected.
- Light rainfall on 28th and 30th July and moderate rainfall on rest of the days is expected. Next five days total rainfall of 81.0 mm is expected.
- Wind mainly from South Easterly, Southerly, South westerly, Westerly direction with 5-8 km/hr. speed is expected.
- Maximum relative humidity is expected to be around 80% - 87% and minimum relative humidity is expected to be around 51% - 72%.
- Maximum temperature is expected to be around 29° - 32°C and minimum temperature is expected to be around 24° - 26° C.

Agrometeorological Advisory

Crop	Stage	Advisory
Aman Rice	Seedling	The transplanting to be continued. Many weeds of grass and sedge family may grow in the rice field. To control the weeds, herbicides like anilophos @ 1.2 kg or Butachlore @ 3 kg or oxydiogen @ 1 kg per ha, mixed with sand, should be applied 4 days after transplanting.
Jute	Mature	Complete the harvesting operation of jute of four months old. At the time of harvesting, cut the leaves in the main-field itself and afterward mix the leaves with top soil by ploughing. This will help in nutrient supply to the next crop. In case of jute-plant grown for seed and late sown crop, this time attack of Bihar hairy caterpillar and semilooper may be prevalent. As suggested in the last week, apply 1.5 ml Endosulphan per lt of water for both the insects.
Cucurbits	-----	In present weather condition attack of powdery mildew may be seen in cucurbits. To control this, spray Dinocap or Trimodorp @ 1 ml/ lt of water. Spray should be done in non rainy day
Chili	-----	The present weather conditions are favorable for occurrence of thrips and mite in chilies. To control spray diphenituran @ 1.25 g./lt. of water is recommended on a non-rainy day.