

Agro-met Advisory Service bulletin for the District South 24 Parganas (Period to 1st August to 5th August,2018) Issued jointly by GKMS, Kakdwip, BCKV and IMD



Bulletin No.:60/2018						
Weather for last four days (28 th July to 30 th July, 2018)						
27/07/18	28/07/18	29/07/18	30/07/18			
0.5	17.7	0.7	0.0			
31.5	32.0	33.5	31.5			
25.5	26.0	26.5	27.0			
92	92	85	82			
0	0	0	0			
Weather for the preceding week (24 th July to 30 th July, 2018)						
	170.4					
Maximum Temperature (°C)			29.5-33.5			
Minimum Temperature (°C)		24.5-27.0				
Relative Humidity (%)		82-96				
Wind Speed (Km ph)			0-1			
	27/07/18 0.5 31.5 25.5 92 0 e preceding v C) C)	four days (28 th July to 30 th 27/07/18 28/07/18 0.5 17.7 31.5 32.0 25.5 26.0 92 92 0 0 e preceding week (24 th July C)	$\begin{array}{ c c c c c c c c c c c c c c c c c c c$			

Weather forecast for next five days (1st August to 5th August,2018)

- Heavy rainfall is predicted on 1st August and light to moderate rainfall is predicted in the remaining days.
- Sky will be overcast in the next five days.
- Wind speed will be 3-10 km/hr. and the predominant wind direction will be Northeasterly to Northwesterly.
- Maximum temperature is expected to be around 31.0° C - 33.0° Cand minimum temperature is likely to be around 26.0° C- 27.0° C.
- ➤ Maximum and minimum relative humidity will be in the range of 87% -91% and 58% -71%.

Weather	based	Agro-A	dvisories
vvcaulti	Dascu	Agi U-A	u v 1501 105

Сгор	Stage	Disease pest	Advisory		
Aman wet	Complete transplanting within first week of August				
seeded paddy	Where field preparation is continuing and fields are found flooded, wait 2-3 days for drain out of				
	excess water then again start transplanting				
Betel vine	Green leaf	Chitla and Rotting			
		green leaf due to	For avoiding rotten of leaf wash the mud at the lower		
		muddy condition	1 0		
		heavy rain	Cover the boundary of garden by polythene		
			For chital disease apply Bordeaux mixture at 2-3 days		
			interval at afternoon hours of sunny days		
Tomato, brinjal,	Vegetative	Wilting due to sur	•		
okra (Early	stage	days after heavy r	ainApply any fungicide at the base during sunny days		
planted)					
Cattle	All stage	Black quarter, for			
		and mouth disease			
		worm infection	medicine at the side of wound		
			Wash the wound of foot and mouth disease with		
			potassium per manganet solution		
			Keep the shade clean and dry for better milk		
	A 11 .		production		
Poultry	All stages	Fungal attack	For preventing fungal attack apply lime		
			For avoiding damp condition in liter apply old news		
			paper and keep the thickness of liter one to one and half		
			inch.		