


Agro-met Advisory Service bulletin for the District

South 24 Parganas

(Period 19thAugust to 23rdAugust, 2017)

Issued jointly by GKMS, Kakdwip, BCKV and IMD



venture for tast times days (15 magast to 17 magast 2017)							
	15/08/17	16/08/17	7/08/17				
Rainfall (mm)	0.3	21.2	0.0				
Maximum Temperature (°C)	34.0	31.0	33.5				
Minimum Temperature (°C)	26.5	26.5	26.0				
Relative Humidity (%)	88	89	88				
Wind Speed (Km ph)							

Weather for the preceding week (11thAugust to 17thAugust,2017)

Rainfall (mm)	44.8
Maximum Temperature (°C)	31.0-34.0
Minimum Temperature (°C)	25.5-29.0
Relative Humidity (%)	85-96
Wind Speed (Km ph)	ı

Weather forecast for next five days (19th August to 23rd August, 2017)

- Light to moderate rainfall is predicted on next five days.
- > Sky will be partly to mainly cloudy the next five days.
- ➤ Wind speed will be 9-13 km/hr. and the predominant wind direction will be Easterly to Southeasterly.
- ➤ Maximum temperature is expected to be around 31.0°C-33.0°C and minimum temperature is likely to be around 26.0°C-27.0°C.
- Maximum and minimum relative humidity will be in the range of 86-88% and 62%-73%.

Weather based Agro-Advisories

Crop	Stage	Disease pest		Advisory		
Paddy aman	Complete transplanting of main field with old seedlings and give first top dressing just					
	after 10 days.					
	Where field is completely destroyed cultivate after early spinach, radish, amaranthus,					
	sweet gourd, pumkin etc.					
Aman paddy (no flood condition)	Start tillering	Apply first top dressing by 12kg urea per bighaMaintain proper water levelCheck the field during morning for any disease or pest attack				
Betel vine	Green leaf	Rotting of leaves near ground level		Drain out excess waterArrange proper air movement inside gardenApply Trichoderma Viridi @5g/liter of water at 15 days interval for two times		
Vadu Kalai	Sowing			Sow seeds@4-5kg/bigha after seed treatment with thirum@3g/kg of seed		
Poultry	Mature bird	Fungal attack	Re Ke Fo	or preventing fungal attack apply lime earrange the litter with lime eep thickness of litter 1inch but never exceed 1 ^{1/2} inch or avoiding damp condition in liter apply old news er evoid wet food use only dry food for feeding purpose		

