

Agro-met Advisory Service bulletin for the District South 24 Parganas (Period to 22nd August to 26th August,2018) Issued jointly by GKMS, Kakdwip, BCKV and IMD



$\mathbf{D}_{\mathbf{v}} = \mathbf{U}_{\mathbf{v}} \mathbf{V}_{\mathbf{v}} $,		to. 21st A	at 2019		
Bulletin No.:66/2018	41-	Date: 21 st August, 2018				
Weather for last for	ur days (17 ^m /	August to 20 th August, 2018)				
	17/08/18	18/08/18	19/08/18	20/08/18		
Rainfall (mm)	23.0	5.4	15.6	36.9		
Maximum Temperature (°C)	33.0	34.0	33.0	34.0		
Minimum Temperature (^o C)	26.0	27.0	27.5	26.5		
Relative Humidity (%)	92	85	89	96		
Wind Speed (Km ph)	0	0	0	1		
Weather for the preceding week (14 th August to 20 th August, 2018)						
Rainfall (mm)		101.8				
Maximum Temperature (°C)		33.0-34.5				
Minimum Temperature (°C)		26.0-27.5				
Relative Humidity (%)		85-96				
Wind Speed (Km ph)		0-1				
	\mathbf{W}_{2} = 4 \mathbf{h}_{2} = \mathbf{f}_{2} = \mathbf					

Weather forecast for next five days (22nd August to 26thAugust,2018)

- > Light to moderate rainfall is predicted in the remaining days.
- Sky will be mainly cloudy in the next five days.
- Wind speed will be 6-16 km/hr. and the predominant wind direction will be South-Easterly to Westerly.
- Maximum temperature is expected to be around 31.0° C - 33.0° Cand minimum temperature is likely to be around 26.0° C- 27.0° C.
- ➤ Maximum and minimum relative humidity will be in the range of 87% -91% and 63% -73%.

Weather	based	Agro-Ad	lvisories
vvcaulti	Dascu	Agro-Au	111301103

Сгор	Stage	Disease pest	Advisory	
Aman paddy	Tillering	Chance of	Maintain proper water level in the field	
		stem borrer	Weeding should be done at regular interval by hand weeding	
			in the main field	
			First top dressing should be done	
			For stem borer attack apply apply granular pesticide	
			Cloranthratrol 0.4g@1.25kg per bigha	
Betel vine	Growing	Chitla due toArrange		ange proper air movement for avoiding dump condition
	leaf	high humidity	ity inside the garden	
			App	bly copper fungicide for chital disease attack
Brinjal,	Flowering to	Rotting and borer		Drenching the soil around seedlings with Diethane
Tomato, Chilli	fruit bearing	pest attack		<u>M-45@2.5g/liter</u> of water for preventing wilting in
				sunny days
				Apply systemic pesticide with sticker
Poultry	Growers and	Fungal attack		Rearrange the litter with new one
	layers			Apply lime on litter
				Avoid wet food only feed them dry food
				The thickness of litter should not be exceeded $1^{1/2}$
				inch

*From NDVI data it is found rainy season in the Southern Part of West Bengal and from SPI data wet condition was found