

Agro-met Advisory Service bulletin for the District Purba Medinipur (Period to 9th May to 13th May,2018) Issued jointly by GKMS, Kakdwip, BCKV and IMD



Bulletin No.:37/2018	Date: 8 th May, 2018					
Weather for last four days (4 th May to 7 th May, 2018)						
	04/05/18	05/05/18	06/05/18	07/05/18		
Rainfall (mm)	0.0	0.0	0.1	0.0		
Maximum Temperature (°C)	32.0	32.5	33.0	34.0		
Minimum Temperature (°C)	24.0	25.0	26.0	28.0		
Relative Humidity (%)	72	80	85	87		
Wind Speed (Km ph)						
Weather for the preceding week (1 st May to 7 th May, 2018)						
Rainfall (mm)	10.1					
Maximum Temperature (°C)		32.0-34.0				
Minimum Temperature (°C)		20.0-28.0				
Relative Humidity (%)		65-87				
Wind Speed (Km ph)	i doth a c					

- Weather forecast for next five days (9th May to 13th May,2018)
 Light rainfall is predicted on 10th May to 11th May and no rainfall is predicted in the remaining days.
- \blacktriangleright Sky will be clear in the next five days.
- > Wind speed will be 13-19 km/hr. and the predominant wind direction will be Southerly to Southwesterly.
- \blacktriangleright Maximum temperature is expected to be around 34.0°C-35.0°C and minimum temperature is likely to be around 26.0° C-27.0°C.
- ➤ Maximum and minimum relative humidity will be in the range of 75% -90% and 43% -56%.

Weather based Agro-Advisories

Crop	Stage	Disease pest	Advisory		
Crop	Stage	Disease pest		Auvisory	
Green	Follow the practice of deep summer ploughing in harvested boro paddy field				
manure	Collect the seed of dhaincha and sow the seed @3-4kg/bigha				
	Prepare land by deep ploughing for cultivation of direct seeded paddy and collect the seeds from				
	certified agency				
Turmeric,	Prepare land for planting of ginger, turmeric etc				
ginger					
Brinjal, Okra	Fruiting	shoot and fruit	borer	To control shoot and fruit borer collect the affecte	
		attack		fruits and buried into the soil	
				If infestation found high then spray Spinosad 48EC	
				@1ml/liter of water.	
Green gram	2 nd pod maturity	Spodoptera a	ttack	Start picking of mature pod of 2 nd flowering during	
	continuing			morning hours	
				For spodoptera attack spray B.T. @1g/liter of water	
				or Spinosad @ 1.5ml/liter of water at afternoon hours	
Guava,citrus	Bend the branch of guava at ground level for better fruiting				
,mango	To prevent fruit drop of citrus spray 10ppm solution of 2,4D				
	To prevent fruit drop of mango spray 15ppm NAA				