

### **Agro-met Advisory Service bulletin for the District**

## Purba Medinipur

# (Period to 19<sup>th</sup> May to 23<sup>rd</sup> May,2018)

### Issued jointly by GKMS, Kakdwip, BCKV and IMD

**Date:** 18<sup>th</sup> May, 2018 Weather for last three days (15<sup>th</sup> May to 17<sup>th</sup> May, 2018)

| ( cather for last three adjs (ie way to 1, way 2010) |          |          |          |  |  |  |
|--|----------|----------|----------|--|--|--|
|  | 15/05/18 | 16/05/18 | 17/05/18 |  |  |  |
| Rainfall (mm)  | 0.0      | 0.0      | 23.5     |  |  |  |
| Maximum Temperature ( °C )                           | 32.0     | 33.0     | 34.0     |  |  |  |
| Minimum Temperature ( °C )                           | 21.0     | 25.0     | 28.0     |  |  |  |
| Relative Humidity (%)                                | 60       | 90       | 95       |  |  |  |

Weather for the preceding week (11<sup>th</sup>May to 17<sup>th</sup> May, 2018)

| Rainfall (mm)              | 65.8      |
|----------------------------|-----------|
| Maximum Temperature (°C)   | 32.0-35.0 |
| Minimum Temperature ( °C ) | 21.0-29.0 |
| Relative Humidity ( % )    | 60-95     |
| Wind Speed ( Km ph )       |           |

#### Weather forecast for next five days (19<sup>th</sup> May to 23<sup>rd</sup> May,2018)

- No rainfall is predicted on 19<sup>th</sup> May and light to moderate rainfall is predicted in the remaining days.
- > Sky will be partly to mainly cloudy in the next five days.

Wind Speed (Km ph)

**Bulletin No.:40/2018** 

- ➤ Wind speed will be 17-20 km/hr. and the predominant wind direction will be Southwesterly.
- Maximum temperature is expected to be around  $33.0^{\circ}$ C  $-34.0^{\circ}$ Cand minimum temperature is likely to be around  $26.0^{\circ}$ C- $27.0^{\circ}$ C.
- Maximum and minimum relative humidity will be in the range of 84% -87% and 57% -62%.

Weather based Agro-Advisories

| weather based Agro-Advisories |  |                        |  |   |  |  |
|-------------------------------|--|------------------------|--|---|--|--|
| Crop                          | Stage  | Disease pest           |  | Advisory  |  |  |
| Dry seeded paddy              | Start sowing of dry seeded long duration paddy after proper tillage operation like ploughing, weeding etcBefore sowing seed should be treated by Diethane M-45 properlyStart to collect seed of wet seeded rice from SSC, NSC etcCollect some variety like Sabita Foundation, Swarna sub-1 certified, IR-64, Hira T.L etc. |                        |  |   |  |  |
| Betel vine                    | Green leaf   | Effect of Nor          |  | Prepare strong fencing for protect the crop from strong windFor better leaf development apply mustard cake and Tasko-5If water stagnation take place then drain out the water immediately |  |  |
| Mango                         | Start to ripe  | Anthracnose fruit crac |  | For anthracnose spray Thiodophenol Methyl@1g/liter of water before 1 month of harvestUse Ethrel for artificial ripening of mango and avoir Carbaid  |  |  |
| Cattle                        | Feed the cattle at morning and evening timeAvoid mung mash and mustard cake in their food and supply arhar or black gram mash and sesame or linseed cakeAvoid grazing during afternoon hours for preventing them from lightning due to Norwester   |                        |  |   |  |  |
| Summer vegetables             | No need of irrigationIf water stagnation found then only drain out the excess water  |                        |  |   |  |  |

<sup>\*</sup>Moderate agriculture vigour was found from NDVI date and mild wet condition was found from SPI data as served from IMD.

