

Agro-met Advisory Service bulletin for the District Purba Medinipur (Period to 1st August to 5th August,2018) Issued jointly by GKMS, Kakdwip, BCKV and IMD



Bulletin No.:60/2018	Date: 31 st July, 2018					
Weather for last four days (28 th July to 30 th July, 2018)						
	27/07/18	28/07/18	29/07/18	30/07/18		
Rainfall (mm)	20.9	46.8	1.2	9.6		
Maximum Temperature (°C)	30.0	31.0	32.0	33.0		
Minimum Temperature (°C)	23.0	24.0	25.0	27.0		
Relative Humidity (%)	75	80	95	98		
Wind Speed (Km ph)						
Weather for the preceding week (24 th July to 30 th July, 2018)						
Rainfall (mm)		112.53				
Maximum Temperature (°C)		29.0-33.0				
Minimum Temperature (°C)		23.0-28.0				
Relative Humidity (%)		65-98				
Wind Speed (Km ph)						

Weather forecast for next five days (1st August to 5th August,2018)

- \blacktriangleright Heavy rainfall is predicted on 1st August and light to moderate rainfall is predicted in the remaining days.
- Sky will be overcast in the next five days.
- Wind speed will be 4-13 km/hr. and the predominant wind direction will be Southwesterly to Northwesterly.
- Maximum temperature is expected to be around 32.0° C -34.0^oCand minimum temperature is likely to be around 26.0° C-28.0^oC.
- ➤ Maximum and minimum relative humidity will be in the range of 84% -88% and 63% -77%.

Weather	based	Agro-A	dvisories
vvcaulti	Dascu	Agi U-A	u v 1501 105

Сгор	Stage	Disease pest	Advisory	
Aman wet	Complete transplanting within first week of August			
seeded paddy	Where field preparation is continuing and fields are found flooded, wait 2-3 days for drain out of			
	excess water then again start transplanting			
Betel vine	Green leaf	Chitla and Rotting of		
		green leaf due to	For avoiding rotten of leaf wash the mud at the lower	
		muddy condition fo	1 0	
		heavy rain	Cover the boundary of garden by polythene	
			For chital disease apply Bordeaux mixture at 2-3 days	
			interval at afternoon hours of sunny days	
Tomato, brinjal,	Vegetative	Wilting due to sunn		
okra (Early	stage	days after heavy rai	nApply any fungicide at the base during sunny days	
planted)				
Cattle	All stage	Black quarter, foot	If the cattle already attacked by then promptly give	
		and mouth disease,	penicillin and Tetracyclin and inoculated these	
		worm infection	medicine at the side of wound	
			Wash the wound of foot and mouth disease with	
			potassium per manganet solution	
			Keep the shade clean and dry for better milk	
			production	
Poultry	All stages	Fungal attack	For preventing fungal attack apply lime	
			For avoiding damp condition in liter apply old news	
			paper and keep the thickness of liter one to one and half	
			inch.	