BIDHAN CHANDRA KRISHI VISWAVIDYALAYA MEDIUM-RANGE WEATHER FORECAST AND AGRO-ADVISORY (Agromet Advisory Services, IMD, MoES)

Directorate of Research, B.C.K.V., Kalyani.



 $E\text{-mail: } \underline{aaskalyani@gmail.com} \ , \\ \underline{bhowmickmahasweta2012@gmail.com} \ \\$

From Kalyani (Nadia, WB)

Weather Forecast for North 24-pgs District

FORECAST PERIOD: : : 22nd August to 26th August, 2018

Weather forecast for next five day

- ➤ Mainly cloudy sky is expected during next 5 days.
- ➤ Light to medium rainfall is expected during next few days, total will be 81 mm.
- ➤ Wind flow mainly from SE and S direction with 6-13 km/hr. speed during next 5 days is expected.
- ➤ Maximum relative humidity is expected to be around 88%-92% and minimum relative humidity is expected to be around 64% 72%.
- Maximum temperature is expected to be around 31°C -33°C and minimum temperature is expected to be around 26°C 27°C .

Agrometeorological Advisory

According to NDVI data sent by IMD agricultural vigour of New Alluvial Zone of West Bengal will be good in the coming week.

Crop	Stage	Advisory
Corn	Sowing	Farmers are advised for sowing of sweet corn & baby corn in raised bed.
Aman	Vegetative	At this time paddy crop is in vegetative stage. Farmers are advised to monitor crop regularly, if population of leaf folder or stem borer is above ETL than broadcast of Cartap 4% granules or carbofuran 3% G @ 10 kg/acre or chlorpyriphos 20 EC @ 2 ml/ litre of water for effective control is recommended. In present weather condition, constant monitoring for brown plant hopper in paddy field is advised If insect population is above ETL spray of Imidacloprid 17.8 % SC@ 1.0 ml / 3 lit of water or is recommended.
Bhindi, Brinjal ,Chilly	Vegetative	Constant monitoring of ladyfinger, brinjal and chilli crop against attack of mite and jassid is advised. To control mite, spraying of phosmite @ 2 ml per litre of water is advised while spraying of Dimethoate 30 EC @ 2 ml/liter of water is advised against jassid when sky remain clear.
Cabbage,	Vegetative	Farmers are advised to monitor borer insect. If insect population is above ETL than, spraying of Spinosad 1.0 ml/4 litre of water is advised

Caulifl ower		when sky is clear against fruit borer, diamond back moth etc.
Chrysant hemum & Marigold	Sowing	Farmers are advised to prepare marigold & chrysanthemum nursery in net house. Proper drainage should be done in the fields.