



Gramin Krishi Mausam Sewa
District Level Agromet Advisory Bulletin
Bidhan Chandra Krishi Viswavidyalaya
 Mohanpur, Dist- Nadia, West Bengal



Agromet Advisory Bulletin

Date : 14-12-2021

Weather Forecast of HOWRAH(West Bengal) Issued On : 2021-12-14(Valid Till 08:30 IST of the next 5 days)

Parameter	2021-12-15	2021-12-16	2021-12-17	2021-12-18	2021-12-19
Rainfall(mm)	0	0	0	0	0
Tmax(°C)	25.1	24.9	25.5	25.1	25.6
Tmin(°C)	14.3	14.0	14.0	13.6	14.8
RH-I(%)	51	55	50	46	55
RH-II(%)	30	27	30	25	24
Wind Speed(kmph)	9.3	10.6	10.4	10.6	11.9
Wind Direction(Degree)	340	338	338	338	326
Cloud Cover(Octa)	5	4	1	0	0

Weather Summary/Alert:

No rainfall is likely to occur during next few days. Partly cloudy to clear sky is expected on next few days . Wind speed is expected to be 9-12 km / hour. Max. RH is expected to be 46% to 55%. Maximum temperature is expected to be around 25.0-26.3 deg C and minimum temperature is expected to be around 13.8-15.5 deg C.

General Advisory:

Prepare drainage channels in the fields to immediately drain out the stagnant water from standing crop which is created due to light / moderate/ heavy rainfall of last week's depression rainfall . Apply Blitox @ 4 g/ lt of water in a clear , sunny weather in the morning or afternoon hour. Weeding is recommended with pre emergence herbicide . Stop applying N fertilizer as top dressing at this moment. Download Mausam, Meghdoot & Damini apps at your smart phone .

SMS Advisory:

Direct seeded spinach , radish , carrot etc may be sown as 2.5 months crop in the fields where newly sown Potato or Mustard are totally affected due to water logging condition for recent depression.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
LENTIL	Drain the excess water immediately . For leaf spot , leaf blight, root rot , wilt etc spray Carbendazim (1 g / lt water) and Tebuconazole (1.5 ml per lt of water) may be applied at 1 week interval.

Crop(Varieties)	Crop Specific Advisory
MUSTARD	Drain the excess water immediately to avoid water logging . For alternaria blight , stem or root rot spray with Metalaxyl and Mancozeb @ 2.5 g /lt of water.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
SPINACH	Direct seeded spinach , radish , carrot etc may be sown as 2-2.5 months crop in the fields where newly sown Potato or Mustard are totally affected due to water logging condition for recent depression rainfall.
POTATO	Drain the excess water immediately. Prophylactic spraying with Mancozeb (for early blight) @ 2.5gm/lt water and Metribuzine @ 10 g/ per drum water (for weed) is recommended in a clear sunny day .