



Gramin Krishi Mausam Sewa

District Level Agromet Advisory Bulletin
Bidhan Chandra Krishi Viswavidyalaya
Mohanpur, Dist- Nadia, West Bengal



Agromet Advisory Bulletin

Date : 31-12-2021

Weather Forecast of 24-PARAGANAS-SOUTH(West Bengal) Issued On : 2021-12-31(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-01-01	2022-01-02	2022-01-03	2022-01-04	2022-01-05
Rainfall(mm)	NA	NA	NA	NA	NA
Tmax(°C)	NA	NA	NA	NA	NA
Tmin(°C)	NA	NA	NA	NA	NA
RH-I(%)	NA	NA	NA	NA	NA
RH-II(%)	NA	NA	NA	NA	NA
Wind Speed(kmph)	NA	NA	NA	NA	NA
Wind Direction(Degree)	NA	NA	NA	NA	NA
Cloud cover(octa)	NA	NA	NA	NA	NA

Weather Summary/Alert:

• No rainfall is predicted in next 5 days. • Sky will be clear in next 5 days. • Wind speed will be 13.8-15.4 km/hr and the predominant wind direction will be Northerly . • Maximum temperature is expected to be around 25.1-26.9 degree and minimum temperature is likely to be 13-14.1 degree. • Maximum and minimum relative humidity will be in the range of 50-61% and 28-40%

General Advisory:

As per ERFs deficit rainfall was recorded from 9.12.21 to 15.12.21 and no rainfall was recorded and from 16.12.21 to 22.12.21. Based on MME forecast in next two week from 24.12.21 to 06.01.22 above normal rainfall is forecasted. Based on MME forecast Tmax will be mostly below normal in Week 1 and 2. and for Tmin, it will show mostly below normal in Week 1 and week 2

SMS Advisory:

As Flowering of lathyrus start spray 2% DAP for better flowering

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
-----------------	------------------------

Crop(Varieties)	Crop Specific Advisory
RICE	<p>Boro paddy Early sown boro paddy with one month age • Where seedlings are more than one month age prepare main field for transplanting • After drained out of water from seed bed apply urea before 7 days of transplanting so that seedlings become strong and never bend down in main field due to strong Northern wind • During land preparation of main field follow basal dose of fertilizer as urea@9kg/bigha, SSP@54kg/bigha and MOP@11kg/bigha • During land preparation apply 0.66 Tone compost • As temperature is low maintain minimum water level during transplanting. For nursery bed with growing seedlings (15 DAS) • Chance of Root rot and yellowing of seedlings due to low temperature • Make shade by transparent polythene over seed bed • Follow water treatment (fill the bed with water during evening and remove the water at morning hours) for temperature conservation when seedlings are already grown up • Stirring the seedlings for removing fog over seedlings. • Apply boron and zinc EDTA 0.5g/liter of water for getting healthy seedlings • Apply Diethane M-45@2.5g/liter of water after complete drain out of water from seed bed with growing seed lings • Avoid urea application on growing seed bed for avoiding its cooling effect.</p>

Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
CHICK	<p>Poultry Chicks, Layers and growers Effect of extreme low temperature • In case of new born chicks keep them litter spread over old news paper • Maintain the temperature of brooder house with electric bulb • Where poultry house made over earthen floor use dimmed burning coal throughout the night to maintain the temperature inside chick's place</p>