

Gramin Krishi Mausam Sewa District Level Agromet Advisory Bulletin Bidhan Chandra Krishi Viswavidyalaya Mohanpur, Dist- Nadia, West Bengal



# **Agromet Advisory Bulletin**

Date : 12-04-2022

Weather Forecast of District PASCHIM-BARDHAMAN(West Bengal) Issued On : 2022-04-12(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-04-13	2022-04-14	2022-04-15	2022-04-16	2022-04-17
Rainfall(mm)	0.0	0.0	0.0	0.0	5.0
Tmax(°C)	38.0	39.0	39.0	40.0	39.0
Tmin(°C)	27.0	26.0	27.0	27.0	26.0
RH-I(%)	50	51	27	26	49
RH-II(%)	20	29	17	21	26
Wind Speed(kmph)	10.7	14.9	12.9	9.7	10.8
Wind Direction(Degree)	338	338	333	333	336
Cloud Cover(Octa)	3	2	0	0	0

## Weather Summary/Alert:

No rainfall is likely to occur during next few days except light rainfall on 14.04.2022 & 16.4.2022. Clear to partly cloudy sky is expected on next few days. Wind speed is expected to be 10-23 km / hour. Max. RH is expected to be 49% to 51%. Maximum temperature is expected to be around 38.0-40.0 deg C and minimum temperature is expected to be around 26.0-27.0 deg C.

## **General Advisory:**

Light rainfall is expected on 14.4.2022 & 16 .4.2022 . Spraying , fertilizer application and harvesting may be avoided .

### **SMS Advisory:**

Light rainfall is expected on 14.4.2022 & 16 .4.2022 . Spraying , fertilizer application and harvesting may be avoided .

### Horticulture Specific Advisory:

Horticulture(Varieties)

Horticulture Specific Advisory

District Advisory

Horticulture(Varieties)	Horticulture Specific Advisory			
BRINJAL	Brinjal fruit & shoot borer may be seen at this stage . As soon as the insect is detected , the affected parts should be clipped along with the insect and destroyed . Fruits showing any boring should be picked and destroyed . Weed control is must . Spray with Profenophos + Cypermethrin 1.5ml + Acitamiprid 1gm at 15 days interval. After spraying apply prime pack 1.0-2.6-2.6 liquid fertilizer (5 ml) + Humax 1 ml/lt + 10-26-26 (10 kg/ bigha) + I kg Sagarika dana +5 kg Neemcake as topdressing . One solar light trap per bigha will be useful .			
	Chilli • Fruiting stage • Mite and thrips attack. For mite attack spray Dicofol or Propergite . For thrips attack spray Thiamithoxam .			