District Advisory



Gramin Krishi Mausam Sewa District Level Agromet Advisory Bulletin West Bengal University of Animal & Fishery Sciences Khudiram Bose Sarani, Kolkata



Agromet Advisory Bulletin

Date : 26-04-2022

Weather Forecast of District 24-PARAGANAS-NORTH(West Bengal) Issued On : 2022-04-26(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-04-27	2022-04-28	2022-04-29	2022-04-30	2022-05-01
Rainfall(mm)	0	0	0	0	8
Tmax(°C)	40.7	41.4	41.2	39.4	38.0
Tmin(°C)	27.1	27.0	26.7	26.8	26.8
RH-I(%)	74	82	83	83	83
RH-II(%)	31	37	44	43	46
Wind Speed(kmph)	24.8	26.3	14.5	16.1	16.5
Wind Direction(Degree)	204	201	225	287	248
Cloud Cover(Octa)	0	0	0	0	4

Weather Summary/Alert:

Mainly clear to partly cloudy sky is expected during next five days. Light rainfall is expected during next five days. Expected total amount of rainfall is near about 8 mm. Wind flow mainly from W and SW direction and expected speed may be 14 - 26 km/hrs. Maximum relative humidity is expected to be around 74% - 83% and minimum RH is expected to be around 31% - 46%. Maximum temperature is expected to be around 38 - 41 deg C and Minimum temperature is expected to be around 26 - 27 deg C.

General Advisory:

To control tomato fruit borer use 3 - 4 pheromone trap per acre. For weather updates use 'Meghdoot' application and for Lightning alert updates use 'Damini' application.

SMS Advisory:

To control the white fly and jassid of Vindi, spray Difenthiuron @ 0.75g per litre of water.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
	If saline condition occurs in Rice field then you have to apply Zinc Sulphate @ 3 kg/bigha and 2 kg Boron. And after 1 month the same thing may be applied in same dose to this field.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory		
BRINJAL	In case of Brinjal, to control Fruit and Stem Borer, collect affected plant parts and fruit and burn it. Set up light traps and pheromone traps to monitor the moths and their egg-laying @ $1-2$ /Bigha. Spray B.t formulation 5% WP such as dipel @ 1.5 g / lit of water.		

Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
COW	Please repair the poultry and livestock house before rainy season. Vaccinations are necessary as FMD in Cattle and Goat, P.P.R and Goat pox in case of Goat, R.D and fowl pox in backyard poultry.

Fishieries Specific Advisory:

Fishieries(Varieties)	Fishieries Specific Advisory	
COMMON FISH	To prevent ulcer disease of fish, apply 13 kg lime/1 bigha of pond. Then after 7 days, apply 1.3 kg bleaching powder. To make a new pond, you may cultivate some legume crop like Dhaincha (Sesbania bispinosa) on that place before making the pond.	