1/10/23, 1:46 PM District Advisory



## Gramin Krishi Mausam Sewa District Level Agromet Advisory Bulletin Bidhan Chandra Krishi Viswavidyalaya



Mohanpur, Dist- Nadia, West Bengal

# **Agromet Advisory Bulletin**

Date: 10-01-2023

Weather Forecast of District HOWRAH(West Bengal) Issued On: 2023-01-10(Valid Till 08:30 IST of the next 5 days)

Parameter	2023-01-11	2023-01-12	2023-01-13	2023-01-14	2023-01-15
Rainfall(mm)	0.0	0.0	0.0	0.0	0.0
Tmax(°C)	26.0	26.0	27.0	27.0	26.0
Tmin(°C)	11.0	12.0	13.0	15.0	16.0
RH-I(%)	38	43	50	80	77
RH-II(%)	19	20	23	23	25
Wind Speed(kmph)	9.5	8.7	6.0	7.8	9.3
Wind Direction(Degree)	338	342	326	207	243
Cloud Cover(Octa)	0	0	0	0	0

#### Weather Summary/Alert:

No rainfall is likely to occur during next few days. Clear sky is expected on next few days. Wind speed is expected to be 6-10 km / hour. Max. RH is expected to be 38% to 77%. Minimum RH is expected to be 19%-26%. Maximum temperature is expected to be around 26.0-27.0 deg C and minimum temperature is expected to be around 11.0-16.0 deg C.

### **General Advisory:**

Please download Mausam, Meghdoot & damini app at your smart phone to get weather forecast and agromet advisory.

#### **SMS Advisory:**

Spraying, fertilizer application, harvesting etc may be done in the coming week as no rainfall is predicted.

### **Horticulture Specific Advisory:**

Horticulture(Varieties)	Horticulture Specific Advisory	
РОТАТО	Since Potato is a tuber crop , soil must be loose and friable . Light spading and frequent hand weeding make the soil loose & friable which enhance growth of tuber smoothly . Partly to mainly cloudy sky , foggy morning , maximum temperature 24.8 to 28.8 deg C , minimum temperature 7-10 deg C may cause early blight of Potato in the second or third week of January . To prevent this disease Mancozeb 2.5 g/lt water or Metalaxyl plus Mancozeb 2.5g/ lt water or Copper oxychloride 4g/ lt water spraying may be recommended. Hexaconazole spraying has been highly effective against the disease .	
	For control of Brinjal fruit & shoot borer , farmers are advised to apply Triazophos 40 EC $@$ 1.5 ml or Cartap Hydrochloride $@$ 0.75 g per lt of water .	