



Agromet Advisory Bulletin

Date : 30-01-2024

Weather Forecast of District MEDINIPUR-EAST(West Bengal) Issued On : 2024-01-30(Valid Till 08:30 IST of the next 5 days)

Parameter	2024-01-31	2024-02-01	2024-02-02	2024-02-03	2024-02-04
Rainfall(mm)	0.0	1.0	6.0	3.0	0.0
Tmax(°C)	27.0	26.0	26.0	26.0	27.0
Tmin(°C)	17.0	17.0	17.0	16.0	15.0
RH-I(%)	82	85	90	89	56
RH-II(%)	47	48	51	67	40
Wind Speed(kmph)	15	15	16	12	11
Wind Direction(Degree)	193	180	162	191	22
Cloud Cover(Octa)	1	2	3	1	0

Weather Summary/Alert:

Light rainfall is predicted from 1st to 3rd February. • Sky will be mainly clear in next 5 days . • Wind speed will be 11-15 km/hr and the predominant wind direction will be Northerly to Southerly. • Maximum temperature is expected to be around 26.0-27.0 degree and minimum temperature is likely to be 15.0-17.0degree. • Maximum and minimum relative humidity will be in the range of 56-90% and 40-67%.

General Advisory:

As per extended range forecast no rainfall is recorded from 11.1.24 to 17.1.24 and above normal rainfall is recorded from 18.1.24 to 24.1.24. Based on MME forecast above normal rainfall is predicted from 26.1.24 to 8.2.24. Based on MME forecast Tmax will be above normal in Week 1 and week 2. For Tmin, it will show below normal in Week 1 and week 2.

SMS Advisory:

As light rainfall is predicted and fog is there for late blight attack of potato apply compound fungicide of Metalaxyl and Mancozeb in clear weather.

Crop Specific Advisory:

Crop (Stage)	Crop Specific Advisory
	Boro paddy Transplanting almost complete /vegetative stage • Those who applied urea fertilizer in nursery bed 7-10 days before and the growing seedlings turned light reddish immediate complete transplanting. Maintain shallow depth during transplanting and allow minimum water level in main field after transplanting. • Those who applied urea in nursery bed just before

Crop (Stage)	Crop Specific Advisory		
	2-3 days, start transplanting 3-4 days after as there is a chance of rainfall up to 2nd February. • If transplanted seedlings in main filed found died due to strong northern wind, gap filling should be done with extra seedling. • Apply pre emergence weedicide just after 3-4 days of transplanting.		
LENTIL (Flowering)	 Lentil, lathyrusFlowering stageRust attack found • Apply DAP for better flowering • Spray Chlorothalonyl @2ml per liter of water at afternoon hours. No spraying and fertilizer application should be done from 1st to 3rd February 		

Horticulture Specific Advisory:

Horticulture (Stage)	Horticulture Specific Advisory		
CHILLI (Flowering)	ChilliVegetative to flowering stageLeaf curl due to strong Northern wind. • Spray Imidachlorpit @3ml per 10 liter of wate or Thiamethoxon @ 1g per liter of water. • No spraying and fertilizer application should be done from 1st to 3rd February		
MANGO (Flowering)	MangoFlowering stageFungal and hopper attack found • Spray Diet M-45 @2.5g per liter of water for fungal attack. • Spray Carbaryl @3g p liter of water for hopper attack. • No spraying and fertilizer application s be done 1st to 3rd February		