



**Gramin Krishi Mausam Sewa**  
**District Level Agromet Advisory Bulletin**  
**Bidhan Chandra Krishi Viswavidyalaya**  
 Mohanpur, Dist- Nadia, West Bengal



## Agromet Advisory Bulletin

Date : 05-03-2024

Weather Forecast of District MEDINIPUR-EAST(West Bengal) Issued On : 2024-03-05(Valid Till 08:30 IST of the next 5 days)

Parameter	2024-03-06	2024-03-07	2024-03-08	2024-03-09	2024-03-10
Rainfall(mm)	0.0	0.0	0.0	0.0	0.0
Tmax(°C)	32.0	32.0	32.0	32.0	32.0
Tmin(°C)	20.0	20.0	21.0	21.0	21.0
RH-I(%)	79	67	79	45	38
RH-II(%)	32	28	43	29	24
Wind Speed(kmph)	11	10	8	10	9
Wind Direction(Degree)	248	225	333	22	243
Cloud Cover(Octa)	0	0	3	0	0

### Weather Summary/Alert:

No rainfall is predicted in next 5 days. • Sky will be to mainly clear in next 5 days . • Wind speed will be 8-11 km/hr and the predominant wind direction will be Easterly to North-Westerly. • Maximum temperature is expected to be around 32.0 degree and minimum temperature is likely to be 20.0-21.0 degree. • Maximum and minimum relative humidity will be in the range of 38-79% and 24-43%.

### General Advisory:

As per extended range forecast normal rainfall is recorded from 15.2.24 to 21.2.24 and deficit in rainfall is recorded from 22.2.24 to 28.2.24. Based on MME forecast below normal rainfall is predicted from 1.3.24 to 14.3.24. Based on MME forecast Tmax will be below normal in Week 1 and week 2. For Tmin, it will show belownormal in Week 1 and week 2. • Advisory for orchard mainly for Bauipur and Sonarpur block of South 24 Parganas. • Advisory for paddy and vegetables mainly for all the blocks except Baruipur, Sonarpur and Thakurpukur Maheshtala block of South 24 Parganas. • Advisories for flower mainly for Pashkura block of East Medinipur

### SMS Advisory:

As no rainfall is predicted and temp. is rising, apply irrigation at afternoon hours in betel vine garden and summer vegetables field. Start for harvesting of lathyrus and gram.,

### Crop Specific Advisory:

Crop (Stage)	Crop Specific Advisory
RICE (Tillering)	Boro paddy --Active tillering stage -- Bacterial Blast and fungal blast, brown spot due to high day temperature and low night temperature with morning fog . --Stemborror attack • For blast attack apply a mixture of Streptocyclin @5g per liter of water and Tycyclozole@6g per liter of water. • For brown spot apply trycyclozole@ 5g per 10 liter of water. • Apply the spraying before irrigation when minimum water is there. • As attack is found huge so apraying must be done at an interval of seven days • Apply the fungicide and

Crop (Stage)	Crop Specific Advisory
	bactericide throughout the whole plant. • For stemborer attack spray Propenophos@1.5ml/liter of water

### Horticulture Specific Advisory:

Horticulture (Stage)	Horticulture Specific Advisory
CUCUMBER (Fruit formation)	Cucurbits -- fruiting • Late Blight due to morning fog and rising day temperature • Downey mildew (burning symptom) • Spray during afternoon hours Mancozeb 75% WP@ 2.5g per liter of water for blight attack • Apply Chlorothalonil (Koboch) @2g per liter of water for Downey mildew attack.
JACK FRUIT (Fruit formation)	Jack fruit Tender fruiting stage Blackish of tender fruit and rotten. • Destroy the rotten fruit outside from the garden. • Apply Mancozeb 0.25% or Copper Oxychloride 0.25% at an interval of 21 days. • After harvesting cut the branches and apply Copper Oxychloride 0.25%.
MARIGOLD (Flowering)	Flower • (Marigold) Flowering Blight due to fog and large difference between day and night temp. • Pluck the healthy flowers during morning for marketing • Apply irrigation during afternoon hours • Spray SAF @2.5g/liter of water for Blight attack

### Fisheries Specific Advisory:

Fisheries	Fisheries Specific Advisory
CARP	Fishery Cover of plankton over the pond water • Apply alam (fitkiri) @4.5kg/bigha in pond water